



**SOUTH DAKOTA  
STATE UNIVERSITY**  
College of Agriculture, Food  
and Environmental Sciences

South Dakota State University Extension  
South Dakota Agricultural Experiment Station at SDSU

# 2021 South Dakota Conventional Soybean Variety Trial Results Mount Vernon

Jonathan Kleinjan | SDSU Extension Agronomist

Kevin Kirby | Agricultural Research Manager

Shawn Hawks | Agricultural Research Manager

**Location:** 5 miles north and 3 miles east of Mount Vernon (57363) in Davison County, SD  
(GPS: 43.785839°, -98.205028°)

**Cooperator:** Edinger Brothers

**Soil Type:** Houdek-Prosper loams, 0-2% slopes

**Fertilizer:** none

**Previous crop:** corn

**Tillage:** no till

**Row spacing:** 30 inches

**Seeding Rate:** 150,000/acre

**Herbicide:**

- Burndown 2.9 pt/acre Prowl (pendimethanlin) + 11 oz/acre 2,4-D LV6
- Pre: 4 oz/acre Fierce (flumioxazin + pyroxasulfone) + 6 oz/acre Metribuzin
- Post: 2 pt/acre Basagran (bentazon) + 6 oz/acre Clethodim

**Insecticide:** none

**Date seeded:** 5/11/2021

**Date harvested:** 10/7/2021

SDSU Extension is an equal opportunity provider and employer in accordance with the nondiscrimination policies of South Dakota State University, the South Dakota Board of Regents and the United States Department of Agriculture.

Learn more at [extension.sdstate.edu](https://extension.sdstate.edu).

© 2021, South Dakota Board of Regents



**SOUTH DAKOTA STATE  
UNIVERSITY EXTENSION**

## 2021 South Dakota Conventional Soybean Variety Trial Results Mount Vernon

Table 1. Conventional soybean performance results (average of four replications - **Maturity Group I**) at Mount Vernon, South Dakota.

Variety Information		Agronomic Performance			
Brand	Variety	Maturity Rating	Yield (bu/ ac@13%)	Moisture (%)	Lodging Score (1-5)*
Virtue	V1821	1.8	<b>43.4</b>	12.2	1.0
SD AES	Brookings	1.7	<b>40.8</b>	12.9	1.0
Check	AG17X8	1.7	<b>40.5</b>	12.5	1.0
Viking Seed	O.1718N	1.7	<b>40.4</b>	12.6	1.0
Sevita International	Candor	1.9	39.1	11.3	1.0
Viking Seed	1518N	1.5	38.4	12.9	1.0
Virtue	V1621	1.6	37.2	12.1	1.0
MN AES	MN1807CN	1.8	37.0	11.1	1.0
Brushvale Seed	N1958	1.9	36.9	12.8	1.0
Richland IFC	MK146	1.1	34.9	13.1	1.0
MN AES	MN1806CN	1.8	33.5	12.1	1.0
MN AES	MN1312CN	1.3	30.0	11.5	1.0
<b>Trial Average</b>			37.7	12.2	1.0
<b>LSD (0.05)†</b>			4.0	3.3	-
<b>C.V.‡</b>			7.3	0.6	-

\* Lodging Score (1 = no lodging to 5 = flat on the ground)  
† Yield or moisture value required ( $\geq$ LSD) to determine if varieties are significantly different from one another.  
‡ C.V. is a measure of variability or experimental error, 15% or less is acceptable.



**SOUTH DAKOTA STATE  
UNIVERSITY EXTENSION**

**2021 South Dakota  
Conventional Soybean Variety Trial Results  
Mount Vernon**

Table 2. Conventional soybean performance results (average of four replications - **Maturity Group II**) at Mount Vernon, South Dakota.

Variety Information		Agronomic Performance			
Brand	Variety	Maturity Rating	Yield (bu/ac@13%)	Moisture (%)	Lodging Score (1-5)*
Viking Seed	2155N	2.1	<b>47.7</b>	11.3	1.0
Viking Seed	2418N	2.4	<b>46.5</b>	11.9	1.0
Miller Hybrids	2508	2.5	<b>46.0</b>	11.9	1.0
Viking Seed	2018N	2.0	<b>45.8</b>	12.3	1.0
Miller Hybrids	2018	2.0	<b>45.5</b>	12.3	1.0
Miller Hybrids	2368	2.3	42.5	12.8	1.0
Virtue	V2122	2.1	41.9	12.0	1.0
Richland IFC	MK373	2.0	39.0	10.2	1.0
Brushvale Seed	N2358	2.3	38.6	12.0	1.0
Check	AG17X8	1.7	38.5	12.4	1.0
<b>Trial Average</b>			43.2	11.9	1.0
<b>LSD (0.05)†</b>			2.7	0.3	-
<b>C.V.‡</b>			4.3	1.8	-

\* Lodging Score (1 = no lodging to 5 = flat on the ground)

† Yield or moisture value required ( $\geq$ LSD) to determine if varieties are significantly different from one another.

‡ C.V. is a measure of variability or experimental error, 15% or less is acceptable.